

THE GEOMETRY OF COMPLEX SPACE-TIMES WITH TORSION

by

Giampiero ESPOSITO^{1,2}

¹*International Centre for Theoretical Physics, Strada Costiera 11, 34014 Trieste, Italy*

²*Scuola Internazionale Superiore di Studi Avanzati, Via Beirut 2-4, 34013 Trieste, Italy*

Summary: *The necessary and sufficient condition for the existence of α -surfaces in complex space-time manifolds with nonvanishing torsion is derived. For these manifolds, Lie brackets of vector fields and spinor Ricci identities contain explicitly the effects of torsion. This leads to an integrability condition for α -surfaces which does not involve just the self-dual Weyl spinor, as in complexified general relativity, but also the torsion spinor, in a nonlinear way, and its covariant derivative. A similar result also holds for four-dimensional, smooth real manifolds with a positive-definite metric. Interestingly, a particular solution of the integrability condition is given by right-flat and right-torsion-free space-times.*

Twistor theory was created by Roger Penrose as an approach to the quantum-gravity problem where null surfaces and some particular complex manifolds are regarded as fundamental entities, whereas space-time points are only derived objects, since they might become ill-defined at the Planck length [1,2]. It is by now well-known that the building blocks of classical field theory in Minkowski space-time are the α -planes. In other words, one first takes complexified compactified Minkowski space $CM^\#$, and one then defines α -planes as null two-surfaces, such that the metric vanishes over them, and their null tangent vectors have the two-component-spinor form $\lambda^A \pi^{A'}$, where λ^A is varying and $\pi^{A'}$ is fixed by a well-known differential equation. This definition can be generalized to complex or real Riemannian (i.e. with positive-definite metric) space-times provided the Weyl curvature is anti-self-dual, giving rise to the so-called α -surfaces.

We are here interested in the case of complex space-times with nonvanishing torsion (hereafter referred to as CU_4 space-times). This study appears relevant at least for the following reasons: (i) torsion is a peculiarity of relativistic theories of gravitation; (ii) the gauge theory of the Poincaré group leads to theories with torsion; (iii) theories with torsion are theories of gravity with second-class constraints; (iv)

if torsion is nonvanishing, the occurrence of cosmological singularities can be less generic than in general relativity [3]. We have thus studied the problem: what are the conditions on curvature and torsion for a CU_4 space-time to admit α -surfaces (defined as above) ?

The starting point of our calculation, based on the use of the full U_4 -connection with torsion [2], is the evaluation of the Lie bracket of two vector fields X and Y tangent to a totally null two-surface in CU_4 . By virtue of Frobenius' theorem, this Lie bracket is a linear combination of X and Y :

$$[X, Y] = \varphi X + \rho Y \quad , \quad (1)$$

where φ and ρ are scalar functions. Moreover, using the definition of the torsion tensor $S(X, Y)$, one also has:

$$[X, Y] \equiv \nabla_X Y - \nabla_Y X - 2S(X, Y) \quad . \quad (2)$$

Thus, since in CU_4 models the torsion tensor, antisymmetric in the first two indices, can be expressed spinorially as:

$$S_{ab}{}^c = \chi_{AB}{}^{CC'} \epsilon_{A'B'} + \tilde{\chi}_{A'B'}{}^{CC'} \epsilon_{AB} \quad , \quad (3)$$

where the spinors χ and $\tilde{\chi}$ are symmetric in AB and $A'B'$ respectively, and are *totally independent*, one finds by comparison of Eqs. (1) and (2) that [2]:

$$\pi^{A'} \left(\nabla_{AA'} \pi_{B'} \right) = \xi_A \pi_{B'} - 2\pi^{A'} \pi^{C'} \tilde{\chi}_{A'B'AC'} \quad , \quad (4)$$

where we have set $X^a = \lambda^A \pi^{A'}$, $Y^a = \mu^A \pi^{A'}$. This is the desired necessary and sufficient condition for the field $\pi^{A'}$ to define an α -surface in the presence of torsion [1,2]. We now have to derive the integrability condition for Eq. (4). For this purpose, we operate with $\pi^{B'} \pi^{C'} \nabla_{C'}$ on both sides of Eq. (4), we repeatedly use the Leibniz rule and Eq. (4), and we also use the spinor formula for the Riemann tensor and spinor Ricci identities. The relations we need are respectively [2]:

$$\begin{aligned} R_{abcd} = & \psi_{ABCD} \epsilon_{A'B'} \epsilon_{C'D'} + \tilde{\psi}_{A'B'C'D'} \epsilon_{AB} \epsilon_{CD} \\ & + \Phi_{ABC'D'} \epsilon_{A'B'} \epsilon_{CD} + \tilde{\Phi}_{A'B'CD} \epsilon_{AB} \epsilon_{C'D'} \\ & + \Sigma_{AB} \epsilon_{A'B'} \epsilon_{CD} \epsilon_{C'D'} + \tilde{\Sigma}_{A'B'} \epsilon_{AB} \epsilon_{CD} \epsilon_{C'D'} \\ & + \Lambda \left(\epsilon_{AC} \epsilon_{BD} + \epsilon_{AD} \epsilon_{BC} \right) \epsilon_{A'B'} \epsilon_{C'D'} \\ & + \tilde{\Lambda} \left(\epsilon_{A'C'} \epsilon_{B'D'} + \epsilon_{A'D'} \epsilon_{B'C'} \right) \epsilon_{AB} \epsilon_{CD} \quad , \end{aligned} \quad (5)$$

$$\left[\nabla_{C(A'} \nabla_{B')}^C - 2\tilde{\chi}_{A'B'}^{HH'} \nabla_{HH'} \right] \pi^{C'} = \tilde{\psi}_{A'B'E'}^{C'} \pi^{E'} - 2\tilde{\Lambda} \pi_{(A'} \epsilon_{B')}^{C'} + \tilde{\Sigma}_{A'B'} \pi^{C'} . \quad (6)$$

As usual, the *twiddle* symbol denotes spinor or scalar quantities independent of their untwiddled counterpart. The spinors ψ and $\tilde{\psi}$ are the Weyl spinors, and are thus invariant under conformal rescalings of the metric [1]. The spinors Σ and $\tilde{\Sigma}$ express the antisymmetric part of the Ricci tensor. Using also the well-known property $\lambda_A \lambda^A = \pi_{A'} \pi^{A'} = 0$, the integrability condition for α -surfaces is thus found to be [2]:

$$\begin{aligned} \tilde{\psi}_{A'B'C'D'} = & -4\tilde{\chi}_{A'B'AL'} \tilde{\chi}_{C'}^{L'A}{}_{D'} + 4\tilde{\chi}_{L'B'AC'} \tilde{\chi}_{A'D'}^{AL'} \\ & + 2\nabla_{D'}^A \left(\tilde{\chi}_{A'B'AC'} \right) . \end{aligned} \quad (7)$$

Note that an analogous result holds if the metric is positive-definite rather than complex [2]. A four-manifold with nonvanishing torsion and positive-definite metric is here denoted by RU_4 . Interestingly, a particular solution of Eq. (7) is given by $\tilde{\psi}_{A'B'C'D'} = 0$, $\tilde{\chi}_{A'C'AB'} = 0$. By analogy with complexified general relativity, the particular CU_4 and RU_4 space-times satisfying these additional conditions are here called right-flat and right-torsion-free. This means that the surviving Weyl and torsion spinors, i.e. ψ_{ABCD} and $\chi_{AB}^{CC'}$, do not affect the integrability condition (7) for α -surfaces.

Acknowledgments

I am very grateful to the referees at GRG for correcting a large number of errors affecting the original version of a paper of mine, which I later revised in a substantial way, writing Ref. [2]. The successful completion of my research, motivated by Ref. [2], would have been unconceivable without the substantial help I received from these two anonymous referees. I am also very grateful to Professor Abdus Salam, the International Atomic Energy Agency and UNESCO for hospitality at the ICTP, and to Professor Dennis Sciama for hospitality at the SISSA of Trieste. Last but not least, I thank Paul Haines for finding a misprint in the manuscript.

References

- [1] R. S. WARD, R. O. WELLS, *Twistor Geometry and Field Theory*, Cambridge Monographs on Mathematical Physics. (Cambridge University Press, Cambridge, 1990)
- [2] G. ESPOSITO, *DSF preprint n. 20* (1991) INFN and Dipartimento di Scienze Fisiche di Napoli.
- [3] G. ESPOSITO, *Fortschr. der Physik* **40** (1992) 1-30.